

Contribution to the Knowledge of the Groundwater Communities from Northern Morocco

Mohammed YACOUBI-KHEBIZA 1; Mohammed MESSOULI 1;
Nicole COINEAU 2; Abdelmoumen Fakher el ABIARI 3

1 - Faculté des Sciences Semlalia, Laboratoire d'Hydrobiologie, B.P. 2390, Marrakech, Morocco.

e-mail: yacoubi@ucam.ac.ma and messouli@ucam.ac.ma

2 - Université P. et M. Curie, Paris 6, Observatoire Océanologique de Banyuls, Laboratoire Arago,
F-66650 Banyuls-sur-Mer, France. e-mail: n.coineau@obs-banyuls.fr

3 - Centre National d'Hydrobiologie et de Pisciculture, BP 11, Azrou, Morocco.

e-mail: elabiari@yahoo.fr

Abstract

The groundwater fauna from the Rif region, in northern Morocco, is highly diversified. Several stygobite taxa are new for science: oligochaetes (species of *Trichodrilus*, genus and species of Phallodrilinae), gastropods (new species of *Hadziella*, *Heideella*, *Pseudoislamia* and *Belgrandia*), amphipods (ten new species of *Pseudoniphargus*), and isopods (*Microcharon alamae* has been recently described). Some taxa are reported for the first time from Morocco, i.e. the oligochaetes Phallodrilinae, as well as the crustacean *Metastenasellus* sp. (isopod). As in other Moroccan regions already investigated, the distribution of groundwater communities, especially peracarid crustaceans, is strongly related to water quality, substrate grain size and the local palaeogeography. The latter factor provides an understanding of the observed community differences both between sites and with other Moroccan regions.

Introduction

The Moroccan groundwater fauna was still poorly known until the beginning of the eighties (CHAPPUIS, 1953; BALAZUC & RUFFO, 1953; KARAMAN & PESCE, 1980; PESCE *et al.*, 1981). More systematic stygobiological investigations were recently performed, first in the Marrakesh region, then in other parts of Morocco: Goulmima (BENAZZOZ, 1983), Tiznit (BOULAL, 1988), Guelmim (IDBENNACER, 1990), El Jadida (FAKHER EL ABIARI, 1999) and the Rifian region. These last sampling were performed with the aim of both making more complete the inventory of known stygobiontic taxa from Morocco and testing once more the possible relationships between the biodiversity of the subterranean aquatic fauna and groundwater quality. This paper finally presents the first data on the Rifian stygobiontic fauna.

Study Area and Methods

The Rifian domain is the Moroccan part of the Betico-Rifian arc which is the western end of the Alpine ranges in the Mediterranean Basin (FRIZON DE LA MOTTE *et al.*, 1991). It is formed by several alpine nappes including three main structural units: the Sebides which are the deepest ones, with a base of peridotites, then the Ghomarides which are of Paleozoic, Triassic, Liasic and Eocene age, and finally the Rifian range formed by a pile of folds and small carbonated nappes of the upper Trias and lower Lias periods.

Three sampling campaigns were successively performed in the region. More than sixty wells and ten or so springs were prospected within a range of 300 km, extending from the Atlantic Ocean to the Algerian Boarder (fig.1).

The fauna of wells was generally collected using a phreatobiological net similar to the Cvetkov model, and baited traps, a more efficient method for capturing crawling species (BOUTIN & BOULANOUAR, 1983). These two methods were completed, when possible, by filtering through a plancton net the water extracted by motorized pumps. The fauna of springs was sampled with a net settled at the source when stringing the sediments just upstream from the water emergence (MESSOULI, 1994).

Results

More than 76 taxa were sampled, including 14 species of Oligochaeta, 12 species of Gastropoda and 42 species of Crustacea. The specific richness of the region stygofauna reaches 38 species (representing about 50% of the total community). The most common representatives of the stygofauna are the gastropods (eight species) and the crustaceans (26 species including eight copepods, three ostracods and 15 peracarid and syncarid crustaceans (tab.1).

This global biodiversity is roughly similar to that of other regions of Morocco previously investigated, as 50 species were collected in the Tiznit region (BOULAL, 1988), 38 taxa in the Marrakesh area (BOULANOUAR, 1995) and 59 species in the Guelmim wells, South-West of the Moroccan Anti Atlas (BOUTIN & IDBENNACER, 1989). The Rifian subterranean fauna seems enough diversified when compared with that of Europe and North America (BOTOSANEANU, 1986; JUBERTHIE & DECU, 1994; GIBERT *et al.*, 1994), especially when considering the hydrobiid snails and the peracarid crustaceans.




Figure 1 - Location of the study sites, wells (R) and springs (S), in the Rifian region of Morocco

The Rifian stygofauna is also characterized by a good number of species new for science and by a high rate of endemic species. Five new species of gastropods were collected, including two species also sampled subsequently in the Beni Mellal plain and three species considered as endemic of the Rif, *Giustia rifensis*, *Heideella knidirii* and *Pseudoislamia yacoubii*. Similarly a new endemic genus of Oligochaeta is a representative of the subfamily Phallodrilinae, with a distribution area limited to the North Eastern part of the Rif. Within the crustaceans several endemic species recently collected are still under study. It is the case of three different species of the amphipod *Pseudoniphargus*, of one species of the syncarid *Iberobathynella*, of one species of the stenassellid isopod *Metastenasellus*, and of 3 new cirolanid isopods which include two different species of *Typhlocirolana* and a new genus which will be published as *Rifolana* n.gen. At last one species of the Rifian isopod Microparasellidae was recently described as *Microcharon alamaie* (BOULANOUAR *et al.*, 1997).

From the consideration of this faunistical list some biogeographical remarks are possible. The Oligochaeta Phallodrilinae belong to a marine group and are mentioned for the first time from fresh groundwaters in the Rifian region; it may be considered as a new evidence of the marine origin of a great part of the stygofauna. The occurrence of this representative in the Rif is certainly related to the origin of the Rifian block, derived from the complex Mesozoic and Cenozoic history of the Alboran plate, which emerged and split in the

Mediterranean in a number of blocks which subsequently drifted to North African and Iberian plates. In other respects, the occurrence of species of the amphipod *Metacrangonyx* in this septentrional region is of great biogeographical interest as it extends widely the distribution of a family only known in Morocco from the south of the Idrissid Land.

The amphipods *Pseudoniphargus* now appear as a large group of species occurring also inside the Rifian domain and not only on its southern margin where it was first discovered. Its origin is certainly related (as that of the phreolidrillid *Oligochaeta*) with the history of the Rifian block, especially with the late Tortonian and early Messinian closure of the South-Rifian Channel and the marine transgressions and regressions in the Rifian domain, when the Rifian block collided with the African plate (COINEAU & BOUTIN, 1996, FAKHER *et al.*, 1999). Other Rifian thalassoid stygobiontic groups have representatives in other parts of Morocco, South of the Idrissid Land, and seem related to transgressions and regressions of the Cenomano-Turonian and of the Senonian or Eocene.

Table 1 - List of the aquatic subterranean species in the Rifian region (* : stygobites species, ° : stygophiles species)

Taxa	Taxa	Taxa
Foraminifera	Copepoda	Amphipoda
<i>Ammonia tepida</i> °	<i>Acanthocyclops sp.*</i>	<i>Salentinella sp.*</i>
Oligochaeta	<i>Diacyclops dubbocki</i> °	<i>Pseudoniphargus ruffoi*</i>
<i>Rhyacodrilus sp.</i> °	<i>Diacyclops sp.*</i>	<i>Pseudoniphargus sp.*</i>
<i>Phallogdrilinae n.g. sp.*</i>	<i>Megacyclops viridis</i> °	<i>Pseudoniphargus sp.1*</i>
<i>Trichodrilus sp.*</i>	<i>Metacyclops subdolus*</i>	<i>Pseudoniphargus sp.2*</i>
Gastropoda	<i>Paracyclops imminutus</i> °	<i>Pseudoniphargus sp.3*</i>
<i>Hadziella midarensis*</i>	<i>Paracyclops fimbriatus</i> °	<i>Maghrebidiella sp.*</i>
<i>Heideella knidirii*</i>	<i>Tropocyclops prasinus</i> °	<i>Metacrangonyx sp.*</i>
<i>Heideella andrea*</i>	Ostracoda	Isopoda
<i>Belgrandia sp.</i> °	<i>Pseudocandona albicans</i> °	<i>Proasellus coxalis africanus</i> °
<i>Pseudoislamia rifensis*</i>	<i>Fabaeformiscandona sp1*</i>	<i>Microcharon alamiae*</i>
<i>Mercuria similis</i> °	<i>Fabaeformiscandona sp2</i> °	<i>Typhlocirolana haouzensis aff*</i>
<i>Giustia rifensis*</i>	Syncarida	<i>Typhlocirolana sp.*</i>
<i>Pseudoislamia yacoubii*</i>	<i>Iberobathynella sp.*</i>	<i>Rifolana n.g. sp.*</i>
		<i>Metastenasellus sp.*</i>

Conclusion

The Rifian groundwaters are inhabited by a relatively rich stygobiontic fauna including a number of taxa related to the ancient Mesozoic history of the Rifian block, like in other parts of Morocco, but also by original and generally endemic other taxa related to the more recent Cenozoic history of the Rifian domain. A number of these species, as they were recently collected, are still under study. Before performing a more detailed analysis of the comparative biodiversity of sampled sites, the first results show that the species richness, especially that of stygobionts, is clearly linked with groundwater quality as in other regions of Morocco.

Aknowledgements

This study, and especially the field work, was performed with the support of the Moroccan project "PARS n° 162 Biologie" and of the French-Moroccan project "A.I. of cooperation n°198/SVS/99".

References

- BALAZUC, J., & S. RUFFO. 1953. Due nuove specie del genere *Metacrangonyx* Chevreux delle acque interne del nord africa francese. Mem. Muséo civ. Stor. Nat. Verona, 4 :25-33.
- BENAZZOZ, A. 1983. Recherche de la faune interstitielle dans la vallée de l'Oued Gheris (Versant sud du Haut-Atlas occidental, Maroc). Mem. C.E.A., Univ. Marrakech, 28 p.
- BOTOSANEANU, L. 1986. Stygofauna mundi. A faunistic distributional and ecological synthesis of the world fauna inhabiting subterranean waters (Including the marine interstitia). E. J. Brill/Dr. W. Backhuys, Leiden, 740 p.
- BOULAL, M. 1988. Recherches écologiques sur la faune aquatique des puits de la région de Tiznit (Anti-Atlas occidental, Maroc). Thèse 3ème Cycle Fac. Sci. Marrakech, 228p.
- BOULANOVAR, M. 1995. Faune aquatique des puits et qualité de l'eau dans les régions de Marrakech et des Jbilet. Statut et dynamique d'une population de *Proasellus coxalis africanus* (Crustacés Isopodes) des Jbilet. Thèse Doct. Etat, Fac. Sc. Semlalia, Marrakech, 207 p.
- BOULANOVAR, M., M. YACOUBI-KHEBIZA, C. BOUTIN & N. COINEAU. 1997. Un nouvel Isopode Microparasellide (Crustacea Janiroidea) des eaux souterraines de la région péri-rifaine (Maroc). Bull. Soc. Hist. Nat. Toulouse, 133 : 21-27
- BOUTIN, C., & M. BOULANOVAR. 1983. Méthodes de capture de la faune stygobie: Expérimentation de différents types de pièges appâtés dans les puits de Marrakech. Bull. Fac. Sc. Marrakech, (section Sc. de la vie) 2: 5-21.
- BOUTIN, C., & B. IDBENNACER. 1989. Faune stygobie du sud de l'Anti-Atlas, premiers résultats. Sciences de l'eau, 2(4) :891-904.
- CHAPPUIS, P.A. 1953 . Un nouvel Isopode psammique du Maroc : *Microcerberus remyi*. Vie Milieu, 4(4) : 659-663.
- COINEAU, N., & C. BOUTIN. 1996. Age and origin of *Pseudoniphargus* in Morocco with the description of three new species. Bull. Mus. Civ. St. nat. Verona, 20: 499-516.
- FAKHER EL ABIARI, A. 1999. Rôle des facteurs abiotiques dans la répartition de la faune aquatique souterraine : cas des Crustacés Péracarides stygobies. Mém. Doct. Nat., Univ. Marrakech, 208p.
- FAKHER EL ABIARI, A., Z. OULBAZ, M. MESSOULI & N. COINEAU. 1999. A new species of *Pseudoniphargus* (Crustacea Amphipoda) from subterranean water of North-Eastern Morocco: historical biogeography and evolutionary aspects. Contr Zoology, 68(3): 161-171.
- FRIZON DE LAMOTTE, D., J. ANDRIEUX & J.C. GUEZOU. 1991. Cinématique des chevauchements néogènes dans l'arc bético-rifain : Discussion sur les modèles géodynamiques. Bull. Soc. géol. France, 162(4): 611-625.
- GIBERT, J., D. L. DANIELOPOL & J. STANFORD. 1994. Groundwater Ecology. Academic Press, New York., 571 p.
- IDBENNACER, B. 1990. Recherches écologiques, biogéographiques et démographiques sur la faune aquatique souterraine de la région de Guelmim (Sud-Ouest de l'Anti-Atlas marocain). Thèse de 3ème cycle. Fac. Sc. Semlalia. Marrakech, 275p.
- JUBERTHIE, C., & V. DECU. 1994. Encyclopaedia Biospeologica. Tome 1. Société de Biospéologie. Moulis (France)-Bucarest (Roumanie), 834p.
- KARAMAN, G., & G.P. PESCE. 1980. Researches in Africa by the Zoological Institute of l'Aquila, Italy. V. On three subterranean Amphipods from North Africa (Amphipoda, Gammaridea). Bull. Zool. Museum, Univers Van Amsterdam, 7(20) :197-207.
- MESSOULI, M. 1994. Evolution, Phylogénie et Biogéographie historique des Metacrangonyctidae, Crustacés Amphipodes stygobies du Nord de l'Afrique et des régions voisines. Thèse doct Etat, Fac. Sc. Marrakech, 311p.
- PESCE, G.L., P. TETE & P. DE SIMONE. 1981. Ricerche faunistiche in acque sotterranee freatiche del Maghreb (Tunisia, Algeria, Morocco) et d'ell Egitto. Natura. Soc. Ital. Sci. nat. Museo civ Stor. nat. e acquario civ. Milano.72 (1-2): 63-98.